

## 富山県より発見されたヒゲエリユスリカ属（ Diptera: Chironomidae）の1新種および数種の採 集データ

|     |                                                                                                                                                               |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 著者  | 山本 優                                                                                                                                                          |
| 雑誌名 | 富山市科学博物館研究報告                                                                                                                                                  |
| 号   | 34                                                                                                                                                            |
| ページ | 31-36                                                                                                                                                         |
| 発行年 | 2011-03-15                                                                                                                                                    |
| URL | <a href="http://repo.tsm.toyama.toyama.jp/?action=repository_uri&amp;item_id=947">http://repo.tsm.toyama.toyama.jp/?action=repository_uri&amp;item_id=947</a> |

**A New Species of *Parorthocladius* Thienemann (Diptera: Chironomidae) from Toyama Prefecture, Honshu, Japan, with distributional Data of some chironomid Species**

Masaru Yamamoto  
Kankyou-kagaku Kabushiki-gaisha  
1-7-3, Naka-sakurazuka, Toyonaka, Osaka, 561-0881, Japan

富山県より発見されたヒゲエリユスリカ属 (Diptera: Chironomidae) の1新種および数種の採集データ

山本 優  
環境科学株式会社  
561-0881 大阪府豊中市中桜塚1-7-3

富山市科学博物館の根来尚博士によって富山県内数カ所の雪上で採集されたユスリカを調べる機会を得た。その中に、エリユスリカ亜科 (Orthoclaadiinae), *Parorthocladius* (ヒゲエリユスリカ属) の一新種を見いだすことが出来た。採集者に因んで *negoroi* と命名し記載した。

*Parorthocladius* 属は現在まで、旧北区から3種、東洋区から3種、合計6種が知られている (Lui & Wang 2005)。旧北区3種の内2種は Sasa & Arakawa (1994), Sasa (1996)によりそれぞれ *Hydrobaenus furudoquartus*, *Orthocladius toyamajekeus* の各属の名の下に富山県産の個体に因って日本から記載報告されたものである。これら2種は Sæther et al. (2000)によって *Parorthocladius* に移された。しかし、*Orthocladius toyamajekeus* は模式標本 (NSMT-I-Dipt 4969:国立科学博物館) の調査から *Psectrocladius aquatronus* Sasa, 1979 (NSMT-I-Dipt 4498: 国立科学博物館)の新参シノニムであることが判明した (Yamamoto 2004)。それ故、現時点では *Parorthocladius* は世界に5種が分布することが確認されていることとなる。今回記載報告する *Parorthocladius negoroi* sp. nov.は世界から第6番目の種となる。また、*Parorthocladius negoroi* と同様に積雪時に富山県内から採集された7種を富山県の分布データとして付記した。

キーワード：ユスリカ科；エリユスリカ亜科；ヒゲエリユスリカ属；新種；富山；日本  
Key words: Chironomidae; Orthoclaadiinae; *Parorthocladius*; new species; Toyama; Japan

Introduction

Up to the present, the genus *Parorthocladius* includes six species and has a Palaearctic and Oriental distribution (Lui & Wang 2005). Of these 6 species, two Japanese species, *Hydrobaenus furudoquartus* Sasa & Arakawa and *Orthocladius toyamajekeus* Sasa, were transferred to the genus *Parorthocladius* by Sæther et al. (2000). The latter, however, was treated as a junior synonym of *Psectrocladius aquatronus* Sasa by Yamamoto (2004), based on the investigation of the holotype bearing large pad-like pulvilli. Consequently, up to date, it is recognized that five species are distributed in the world.

By the courtesy of Dr. Hisashi Negoro, Toyama Science Museum, I had an opportunity to examine the 6th species of the genus *Parorthocladius*, together with 7 species; *Diamesa alpina* Tokunaga, *D. japonica* Tokunaga, *Synpotthastia takatensis* (Tokunaga), *Diplocladius cultriger* Kieffer, *Orthocladius frigidus* (Zetterstedt), *O. kanii* Tokunaga and *O. saxosus* Tokunaga. These species were collected from Toyama Prefecture in winter on the snow. Here, I describe the new species and give the distributional data with seven species in the following lines.

## Methods and material

The terminology and measurements mainly follow Sæther (1980). For male hypopygial structure, the paramere and the basal lobe are used as neutral terms to the phallapodeme carrying aedeagal lobe and the inferior volsella of Sæther (1980), respectively. Material used for description and illustration was mounted on a slide in Euparal. The type specimen is deposited in the collections of the Toyama Science Museum, Toyama, Japan.




Fig. 1 *Parorthocladius negoroi* sp. nov., Holotype male imago. A, head, frontal view; B, head, caudal view; C, tentorium, lateral view; D, cibarial pump, ventral view; E, epipharynx, ventral view; F-G, antennal flagellum, F, basal five flagellomeres, G, tip of last flagellomere.

*Parorthocladius negoroi* sp. nov.

Japanese name: Oo-higeeriyusurika (new name)

(Fig. 1 A-G, 2 A-C, 3 A, B)

Type material: Holotype male imago, slide mounted in Euparal: Rendaiji, Nanto-shi, Toyama Pref., Honshu, Japan, 23. I. 2005, Hisashi Negoro (TOYA JI-22940).

Etymology: Named in honour of Dr. Hisashi Negoro, Toyama Science Museum, the collector of the examined material.

Male (n=1)

Total length 3.3 mm. Wing 2.3 mm long, 0.6 mm width; wing length/wing width 3.83.

Coloration. Body including head and legs entirely brown, scutal vittae slightly darkened. Wing hyaline, anterior veins pale brown, posterior veins hyaline. Halter pale brown.

Head (Fig. 1 A-G). First to 5th flagellomeres with 1, 2, 2, 2, 2 blunt sensilla basiconica, respectively; last flagellomere with several pointed sensilla basiconica and ringed sensilla coeloconica. AR 1.40. First to ultimate flagellomere lengths (in  $\mu\text{m}$ ): 64, 24, 24, 20, 24, 24, 24, 28, 28, 28, 28, 32, 536. Palpomere lengths (in  $\mu\text{m}$ ): 36, 52, 112, 104, 148; with 0, 9, 21, 14, 17 setae, respectively; third palpomere with 2 sensilla clavata. Vertex with 17-19 temporal setae, uniserial. Clypeus with 12 setae. Cibarial pump 60  $\mu\text{m}$  long, cornua well developed.

Thorax (Fig. 2 A, B). Antepronotum well developed, nearly parallel-sided dorsally, distinctly divided by v-shaped notch and contacted with scutum through median commissure. Lateral antepronotals 4; dorsocentals 9-11, uniserial; no acrostichal; prealars 5, uniserial; no supraalar. Preepisternu with 1 seta mid-dorsally. Scutellum with 22 setae, biserial.




Fig. 2 *Parorthocladius negoroi* sp. nov., Holotype male imago. A, thorax, dorsal view; B, thorax, lateral view ; C, wing.

Table 1. Lengths (in  $\mu\text{m}$ ) and proportions of legs of *Parorthocladius negoroi* sp. nov., male (n =1).

|    | fe  | ti  | ta1 | ta2 | ta3 | ta4 | ta5 | LR   | BV   | SV   |
|----|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| p1 | 720 | 736 | 608 | 384 | 280 | 200 | 128 | 0.83 | 2.08 | 2.40 |
| p2 | 792 | 712 | 400 | 256 | 208 | 128 | 120 | 0.56 | 2.67 | 3.76 |
| p3 | 856 | 840 | 520 | 328 | 256 | 152 | 128 | 0.62 | 3.90 | 3.26 |

Wing (Fig. 2 C). Anal lobe moderately developed. Costal extension nearly as long as r-m cross-vein,  $40\ \mu\text{m}$  long. VR 1.10. R with 7-8 setae on basal 1/2.  $R_1$  and  $R_{4+5}$  without seta. Brachiolum with 1 median seta; with 10 basal, 3 median and 10 subapical sensilla campaniformia. Squama with 34 setae, biserial.

Legs. Fore-, mid and hind coxae with 5, 4, 6 marginal setae, respectively; fore-, mid and hind trochanters with 7, 4, 5 marginal setae, respectively. Spur of fore tibia  $62\ \mu\text{m}$  long; mid tibia with antero-ventral spur  $30\ \mu\text{m}$  and postero-ventral one  $32\ \mu\text{m}$  long; hind tibia with antero-ventral spur  $22\ \mu\text{m}$  long and postero-ventral one  $66\ \mu\text{m}$  long. Tibial comb of hind leg composed of 11 spine-like setae. Pseudospurs present on 1-2 tarsomeres of mid and hind legs. Lengths and proportions of legs as in table 1.

Hypopygium (Fig. 3 A. B). Anal point gradually tapering to rounded apex,  $58\ \mu\text{m}$  long,  $40\ \mu\text{m}$  wide at base and with 11 short lateral setae. Sternum IX with 8-12 setae, dorsolaterally. Transverse sternapodeme without oral projection, and with anterior margin strongly arched. Paramere  $74\ \mu\text{m}$ . Virga composed of needle-like 2 spines,  $32\ \mu\text{m}$  long. Gonocoxite  $220\ \mu\text{m}$  long. Gonostylus slightly arched,  $108\ \mu\text{m}$  long. Basal lobe double layered, with dorsal part rather wide and rectangular, well sclerotized. Crista dorsalis well developed. Megaseta  $10\ \mu\text{m}$  long.

Distribution. The species is known from type locality only.




Fig. 3 *Parorthocladius negoroi* sp. nov., Holotype male imago, male hypopygium. A, dorsal view; B, ventral view.

Remarks. *Parorthocladius negoroi* sp. nov. resembles *P. nudipennis* (Kieffer) from Europe and Siberia in the hypopygial structure (Brundin, 1956; Cranston et al., 1989), but is easily separable from the latter by the following combination of characters: scutellum with biserially arranged setae, preepisternu with 1 seta, squama with many biserially arranged fringed setae, anal point long and reaching the level of distal end of dorsal part of basal lobe, transverse sternapodeme strongly arched and without oral projection.

Some other species collected at the mountainous region form Toyama Prefecture on the snow  
Diamesinae

1. *Diamesa alpina* Tokunaga, 1936 (Fig. 2, Makarchenko & Yamamoto 1995)

Specimens examined: 1♂, 1♀, Fujibashi, Tateyama-machi, Naka-Niikawa Gun, 18. I. 2004; 3♂♂, as previous except Shichihimedaira, 28. I. 2005; 1♀, as previous except Katsuradai, 2. XII. 2006.

2. *Diamesa japonica* Tokunaga, 1936 (Fig. 4-A)

Specimens examined: 1♂, Fujibashi, Tateyama-machi, Naka-Niikawa Gun, 3. I. 2005; 1♀, Rendaiji, Nanto-shi, 23. I. 2005.




Fig. 4 Male hypopygia. A, *Diamesa japonica* Tokunaga; B, *Symptothastia takatensis* (Tokunaga); C, *Orthocladius saxosus* (Tokunaga).

3. *Synpotthastia takatensis* (Tokunaga, 1964) (Fig. 4-B)

Specimens examined: 1♂, Rendaiji, Nanto-shi, 23. I. 2004.

Orthoclaadiinae

4. *Diploccladius cultriger* Kieffer, 1908 (Fig. 53, Yamamoto 2010)

Specimens examined: 4♂♂, 2♀♀, Fujibashi, Tateyama-machi, Naka-Niikawa Gun, 2. II. 2005.

5. *Orthoccladius frigidus* (Zetterstedt, 1838) (Fig. 63-C, Yamamoto 2010)

Specimens examined: 2♂♂, 2♀♀, Fujibashi, Tateyama-machi, Naka-Niikawa Gun, 18. I. 2004; 1♂, as previous except Nanahime-daira, 28. I. 2005.

6. *Orthoccladius kanii* (Tokunaga, 1939) (Fig. 63-D, Yamamoto 2010)

Specimens examined: 1♂, 1♀, Kanjoji, Nanto-shi, 1. II. 2004; 1♂, Ranjoyama, Tonami-shi, 18. III. 2005.

7. *Orthoccladius saxosus* (Tokunaga, 1939) (Fig. 4-C)

Specimens examined: 1♂, 1♀, Fujibashi, Tateyama-shi, Naka-Niikawa Gun, 3. I. 2005.

Acknowledgements

I am deeply indebted to Dr. Hisashi Negoro, Toyama Science Museum, Toyama, Japan for the material.

References

- Brundin, L., 1956. Zur Systematik der Orthoclaadiinae (Dipt. Chironomidae). Reports from Institute of Freshwater Research, Drottningholm 37: 5-185.
- Cranston, P. S., Oliver, D. R. & O. A. Sæther, 1989. The adult males of Orthoclaadiinae (Diptera: Chironomidae) of the Holarctic region - keys and diagnoses. In Wiederholm, T. (ed.), Chironomidae of the Holarctic region. Keys and diagnoses. Part 3. Adults males. Entomologica Scandinavica Supplement 34: 165-352.
- Liu Y. & Wang X., 2005. A review of the genus Parorthoccladius Thienemann from China (Diptera: Chironomidae: Orthoclaadiinae). Zootaxa. 802: 1-8.
- Makarchenko, E., and Yamamoto, M., 1995. Chironomids of the Diamesinae (Diptera, Chironomidae) from Japan. V. New and little-known species of Diamesa Maigen. Japanese Journal of Entomology, 63 (2): 297-301.
- Sæther, O. A., 1980. A glossary of chironomid morphology terminology (Diptera: Chironomidae). Entomologica Scandinavica Supplement 14: 1-51.
- Sæther, O. A., Ashe, P. and Murray, D. A., 2000. Contribution to a Manual of Palaearctic Diptera. Appendix. A. 6. Family Chironomidae. Pages 113-334, In Papp, L. and Darva, B. (Editors). Science Herald, Budapest: 604pp.
- Sasa, M., 1996. Seasonal distribution of the chironomid species collected with light traps. - At the side of two lakes in the Toyama City Family Park. Toyama Prefectural Environmental Science Research Center, 1996 (March): 15-85.
- Sasa, M. and Arakawa, R., 1994. Studies on the chironomid midges collected in Toyama and other areas of Japan, 1994. Part 3. Seasonal changes of chironomid species emerging from Lake Furudo. Toyama Prefectural Environmental Science Research Center, 1994: 88-109.
- Yamamoto, M., 2004. A catalog of Japanese Orthoclaadiinae (Diptera: Chironomidae). Makunagi/ Acta Dipterologica, (21): 1-121.
- Yamamoto, M., 2010. Chironomidae of Japan -In Nihon Yusurika Kenkyu-kai (Ed.) - IV Orthoclaadiinae: 96-145. (in Japanese).